

IN THE CLAIMS:

The text of all pending claims, (including withdrawn claims) is set forth below. Cancelled and not entered claims are indicated with claim number and status only. The claims as listed below show added text with underlining and deleted text with ~~striketrough~~. The status of each claim is indicated with one of (original), (currently amended), (cancelled), (withdrawn), (new), (previously presented), or (not entered). Please AMEND claims 24 in accordance with the following:

1 (original): A method for producing an aromatic carbonate, comprising:

(1) performing a reaction between an organometal compound and carbon dioxide to obtain a reaction mixture containing a dialkyl carbonate formed by the reaction,

(2) separating said dialkyl carbonate from said reaction mixture to obtain a residual liquid, and

performing the following steps (3) and (4) in either order, or partially or wholly simultaneously:

(3) reacting said residual liquid with an alcohol to form at least one organometal compound and form water and re-

moving said water from said organometal compound, and

(4) reacting said dialkyl carbonate separated in step (2) with an aromatic hydroxy compound to obtain an aromatic carbonate.

2 (original): The method according to claim 1, wherein said aromatic carbonate obtained in step (4) is at least one compound selected from the group consisting of an alkyl aryl carbonate and a diaryl carbonate.

3 (original): The method according to claim 1 or 2, wherein, in step (3), said organometal compound having said water removed therefrom is recycled to step (1).

4 (previously amended): The method according to claim 1 or 2, wherein, in step (4), an alcohol which is generated together with said aromatic carbonate is recycled to step (3).

5 (previously amended): The method according to claim 1 or 2, wherein a dialkyl carbonate recovered in step (4) is recycled to step (4).

6 (previously amended): The method according to claim 1 or 2,

wherein a cycle of steps (1) to (4) is repeated at least one time.

7 (previously amended): The method according to claim 2, wherein said aromatic carbonate obtained in step (4) is an alkyl aryl carbonate and which, after step (4), further comprises the following step (5):

(5) subjecting said alkyl aryl carbonate to a disproportionation reaction to obtain a diaryl carbonate.

8 (original): The method according to claim 7, wherein, in step (5), a dialkyl carbonate which is generated together with said diaryl carbonate is recycled to step (4).

9 (original): The method according to claim 7 or 8, wherein a cycle of steps (1) to (5) is repeated at least one time.

10 (previously amended): The method according to claim 1 or 2, wherein, in step (1), said organometal compound is used in an amount which is 1/200 to 1 time the stoichiometric amount relative to the amount of said carbon dioxide.


11 (previously amended): The method according to claim 1 or 2,

wherein said reaction in step (1) is performed at 20 °C or higher.

12 (previously amended): The method according to claim 1 or 2, wherein said organometal compound used in step (1) is an organometal compound having a metal-oxygen-carbon linkage.

13 (original): The method according to claim 12, wherein said organometal compound having a metal-oxygen-carbon linkage comprises at least one compound selected from the group consisting of:

an organometal compound represented by the formula (1):




wherein:

M^1 represents a metal atom selected from the group consisting of elements belonging to Groups 4 and 14 of the Periodic Table, exclusive of silicon;

each of R^1 and R^2 independently represents a straight chain or branched C_1 - C_{12} alkyl group, a C_5 - C_{12} cycloalkyl group, a straight chain or branched C_2 - C_{12} alkenyl group, a C_7 - C_{20} aralkyl group comprised of unsubstituted or substituted C_6 - C_{19} aryl and alkyl selected from the group consisting of straight chain or branched C_1 - C_{14} alkyl and C_5 - C_{14} cycloalkyl, or an unsubstituted or substituted C_6 - C_{20} aryl group;

each of R^3 and R^4 independently represents a straight chain or branched C_1 - C_{12} alkyl group, a C_5 - C_{12} cycloalkyl group, a straight chain or branched C_2 - C_{12} alkenyl group, or a C_7 - C_{20} aralkyl group comprised of unsubstituted or substituted C_6 - C_{19} aryl and alkyl selected from the group consisting of straight chain or branched C_1 - C_{14} alkyl and C_5 - C_{14} cycloalkyl; and

each of a and b is an integer of from 0 to 2, $a + b = 0$ to 2, each of c and d is an integer of from 0 to 4, and $a + b + c + d = 4$; and
an organometal compound represented by the formula (2):



wherein:

(2)

each of M^2 and M^3 independently represents a metal atom selected from the group consisting of elements belonging to Groups 4 and 14 of the Periodic Table, exclusive of silicon;

each of R^5 , R^6 , R^7 and R^8 independently represents a straight chain or branched C_1 - C_{12} alkyl group, a C_5 - C_{12} cycloalkyl group, a straight chain or branched C_2 - C_{12} alkenyl group, a C_7 - C_{20} aralkyl group comprised of unsubstituted or substituted C_6 - C_{19} aryl and alkyl selected from the group consisting of straight chain or branched C_1 - C_{14} alkyl and C_5 - C_{14} cycloalkyl, or an unsubstituted or substituted C_6 - C_{20} aryl group;

each of R^9 and R^{10} independently represents a straight chain or branched C_1 - C_{12} alkyl group, a C_5 - C_{12} cycloalkyl group, a straight chain or branched C_2 - C_{12} alkenyl group, or a C_7 - C_{20} aralkyl group comprised of unsubstituted or substituted C_6 - C_{19} aryl and alkyl selected from the group consisting of straight chain or branched C_1 - C_{14} alkyl and C_5 - C_{14} cycloalkyl; and

each of e, f, g and h is an integer of from 0 to 2,

$e + f = 0 \text{ to } 2$, $g + h = 0 \text{ to } 2$, each of i and j is an integer of from 1 to 3, $e + f + i = 3$, and $g + h + j = 3$.

14 (previously amended): The method according to claim 1 or 2, wherein said separation of said dialkyl carbonate in step (2) is performed by at least one separation method selected from the group consisting of distillation, extraction and filtration.

15 (original): The method according to claim 14, wherein said separation of said dialkyl carbonate in step (2) is performed by distillation.

16 (original): The method according to claim 15, wherein said separation of said dialkyl carbonate in step (2) is performed by thin film distillation.

17 (previously amended): The method according to claim 1 or 2, wherein said removal of said water in step (3) is performed by membrane separation.

18 (original): The method according to claim 17, wherein

said membrane separation is pervaporation.

19 (previously amended): The method according to claim 1 or 2, wherein said removal of said water in step (3) is performed by distillation.

20 (previously amended): The method according to claim 1 or 2, wherein said alcohol used in step (3) is at least one alcohol selected from the group consisting of an alkyl alcohol having a straight chain or branched C₁-C₁₂ alkyl group, a cycloalkyl alcohol having a C₅-C₁₂ cycloalkyl group, an alkenyl alcohol having a straight chain or branched C₂-C₁₂ alkenyl group, and an aralkyl alcohol having a C₇-C₂₀ aralkyl group comprised of unsubstituted or substituted C₆-C₁₉ aryl and alkyl selected from the group consisting of a straight chain or branched C₁-C₁₄ alkyl and C₅-C₁₄ cycloalkyl.

21 (previously amended): The method according to claim 1 or 2, wherein the alcohol used in step (3) has a boiling point which is higher than the boiling point of water.

22 (original): The method according to claim 21, wherein the alcohol used in step (3) is at least one alcohol selected

from the group consisting of 1-butanol, 2-methyl-1-propanol, an alkyl alcohol having a straight chain or branched C₅-C₁₂ alkyl group, and an alkenyl alcohol having a straight chain or branched C₄-C₁₂ alkenyl group.

23 (previously amended): The method according to claim 21, wherein the alcohol used in step (3) has a boiling point which is lower than that of said aromatic hydroxy compound used in step (4).

24 (currently amended): The method according to claim 13, wherein each of R³ and R⁴ in formula (1) and R⁹ and R¹⁰ in formula (2) independently represents an n-butyl group, a 2-methylpropyl group, a straight chain or branched C₅-C₁₂ alkyl group, or a ~~straight chain or~~ branched C₄-C₁₂ alkenyl group.

25 (previously amended): The method according to claim 1 or 2, wherein, in step (1), said organometal compound is used in at least one form selected from the group consisting of a monomeric form, an oligomeric form, a polymeric form and an associated form.

26 (previously amended): The method according to claim 13,

wherein each of M^1 in formula (1) and M^2 and M^3 in formula (2) represents a tin atom.

27 (previously amended): The method according to claim 1 or 2, wherein said organometal compound used in step (1) is produced from an organotin oxide and an alcohol.

28 (previously amended): The method according to claim 1 or 2, wherein, the amount of said aromatic hydroxy compound used in step (4) is 0.1 to 10,000 times the stoichiometric amount relative to the amount of said dialkyl carbonate used in step (4).

29 (previously amended): The method according to claim 1 or 2, wherein said reaction in step (4) is performed at a temperature in the range of from 50 to 350 °C.

30 (previously amended): The method according to claim 1 or 2, wherein said reaction in step (4) is performed in the presence of a transesterification reaction catalyst.

31 (previously amended): The method according to claim 7 or 8, wherein said reaction in step (5) is performed in the pres-

ence of a disproportionation reaction catalyst.

32 (previously amended): The method according to claim 1 or 2, wherein said aromatic hydroxy compound is represented by the following formula (3):



wherein Ar is a C₅-C₃₀ aromatic group.

33 (original): The method according to claim 32, wherein said aromatic hydroxy compound represented by formula (3) is phenol.

34 (previously amended): The method according to claim 1 or 2, wherein the total content of an aromatic hydroxy compound and a carboxyl group-containing compound in said alcohol used in step (3) is 1,000 ppm or less.

35 (previously amended): An aromatic carbonate produced by the method of claim 1 or 2.

36 (original): A polycarbonate, an isocyanate or a polycarbonate diol produced using the aromatic carbonate of claim 35.

37 (original): The polycarbonate, isocyanate or polycarbonate diol according to claim 36, wherein said aromatic carbonate is a diaryl carbonate.